

ORIGINAL RESEARCH

How frequently are physiological variables recorded in the emergency room in patients with obstructive pulmonary diseases and pneumonia?

Lars K Hægstad^a, Torbjørn Bjøteit^a, Ane Johannessen^b, *Amund Gulsvik^a^a Section of Thoracic Medicine, Institute of Medicine, University of Bergen, Norway^b Centre for Clinical Research, Haukeland University Hospital, Norway

Received 3rd December 2007; revised 7th March 2008; accepted 6th July 2008; online 2nd October 2008

Abstract

Aim: To assess how frequently measurements of relevant physiological variables are made in patients presenting acutely to hospital with obstructive pulmonary diseases (OPD) and pneumonia.

Methods: Patients admitted to three hospitals in western Norway with OPD (asthma or chronic obstructive pulmonary disease (COPD)) and/or pneumonia over a 75-day period were included in the study. The assessing physicians in the emergency room (ER) were asked to measure and record physiological variables according to normal practice.

Results: There were 338 patients (54% women), with an average age of 64 (SD 19) years. The initial diagnoses were OPD (176), pneumonia (103) and a combination of OPD and pneumonia (59). Heart rate and body temperature were recorded in almost all patients, whereas the recording of all other variables varied from 45% to 91%. Most variables were recorded more frequently in OPD than in pneumonia.

Conclusions: The importance of recording respiratory rate, cutaneous oxygen saturation, and peak expiratory flow should be stressed in the training of physicians.

© 2009 General Practice Airways Group. All rights reserved.

LK Hægstad *et al.* *Prim Care Res J* 2009; 18(2): 95-99.

doi:10.3132/pcrj.2008.00057

Keywords asthma, COPD, pneumonia, statistics, hospital, medical education**Introduction**

In textbooks of medicine there is no agreement on which physiological variables should be recorded in patients with suspected chronic obstructive pulmonary disease (COPD) and pneumonia.¹⁻⁶ When acute lung diseases are suspected, physicians frequently measure respiratory rate, heart rate, body temperature and blood pressure.^{7,8} A simple method of assessing lung function in the emergency room (ER) is measurement of peak expiratory flow (PEF),⁹ cutaneous oxygen saturation¹⁰ and/or arterial blood gas concentrations.¹¹ Measurement of C-reactive protein (CRP) may be of value when deciding whether or not a patient presenting with infection should be treated with antibiotics.^{12,13} To our knowledge there is no information on how often

physiological variables are measured and recorded in the ER, and what distinguishes the recordings in OPD from the ones in pneumonia.

The aim of this study was to estimate how frequently respiratory rate, heart rate, body temperature, PEF, cutaneous oxygen saturation, arterial blood gas concentrations, and CRP were recorded in patients admitted to hospital with OPD and/or pneumonia. We wished to establish whether the recording of these variables was associated with the patients' sex, age, initial diagnosis and/or severity of disease.

Methods

All patients admitted with OPD and pneumonia to the Haugesund County Hospital, Haraldsplass Deaconal Hospital

* **Corresponding author:** Institute of Medicine, University of Bergen, Jonas Liesvei 65, 5021 Bergen, Norway
Tel: +47 55 97 32 42 E-mail: amund.gulsvik@med.uib.no

and Haukeland University Hospital were registered between April 1 to June 15 1997.¹⁴⁻¹⁶ The hospital district of Bergen, with Haraldsplass Deaconal Hospital and Haukeland University Hospital, covers 15 municipalities with a total catchment area of 251,936 citizens aged over 15. The Haugesund hospital district, with Haugesund County Hospital, encompasses eleven municipalities with a catchment area of 77,337 citizens aged over 15.¹⁷

In total, 438 patients were admitted with a diagnosis of OPD and/or pneumonia. Physicians and nurses in the ER measured and recorded variables according to normal practice. In addition the physician on duty filled in a registration form for each patient. The form had been tested in a pilot scheme. The patient's age, sex, initial diagnosis and disease severity were recorded. Each receiving physician was asked to classify the patient according to disease severity (i.e. mild, moderate, serious/life threatening) to the best of his or her judgement. There were no demands for objective measurements in order to estimate disease severity – i.e. every physician assessed disease severity based on experience, education and normal hospital routine. The initial diagnosis was made by the referring physician or by the receiving physician if there was no referred diagnosis. The patients were diagnosed with OPD (COPD and asthma), pneumonia or a combination of OPD and pneumonia.

Data were obtained on the frequency of recording of respiratory rate (*V*/minute), peripheral heart rate (*V*/minute), PEF (*L*/minute), body temperature (°C), cutaneous oxygen saturation (%), arterial blood gas concentration (arterial oxygen (P_{aO_2}) and carbon dioxide tension (P_{aCO_2}) (kPa)), and CRP (mg/L).

Heart rate and body temperature were measured by nurses. Respiratory rate was recorded by physicians. Measurement of PEF was conducted by physicians or nurses by means of a simple, portable peak flow meter. The physicians were responsible for measuring CRP and cutaneous oxygen saturation, and obtained blood samples for arterial blood gas analysis; cutaneous oxygen saturation was measured by nurses when asked by physicians, and the patient's venous CRP level was analysed at the local medical biochemistry laboratory.

The registration form was filled in by 26 physicians who were on primary duty at the three hospitals, and 110 nurses measured heart rate, body temperature and cutaneous oxygen saturation. The median number of registered patients per physician was 11 (range 1-23). The data were coded in SPSS ("Statistical Product and Service Solutions) 7.5 for Windows.¹⁵

Statistical analyses

All statistical analyses were performed using SPSS 13.0 for Windows. Pearson's Chi-square test was used to investigate

possible associations between the initial diagnosis and the patients' age, sex and disease severity. We also investigated any association between initial diagnosis and registration of heart rate, body temperature, respiratory rate, PEF, cutaneous oxygen saturation, arterial blood gas concentrations, and CRP by means of Pearson's Chi-square test. Odds ratios with 95% confidence interval were calculated by multiple logistic regression for how frequently these variables were registered depending on sex, age, initial diagnosis and disease severity. The level of significance for all analyses was set at 0.05.

Results

Sex, age, diagnosis and disease severity

The study population was 338 patients (77% of those admitted in the timeframe) with either OPD (176 patients), pneumonia (103 patients), or a combination of the two (59 patients) – see Table 1. Two hundred and twenty-six patients (67%) were admitted to Haukeland University Hospital, 27 (8%) to Haraldsplass Deaconal Hospital and 85 (25%) to Haugesund County Hospital. The average age (SD) was 64 (19), and 54% were women. The patients' ages were, as expected, unevenly distributed, with the median age being 71 years (range 14 – 93 years). There were no differences in age between the sexes and diagnostic groups. Women (63%) were admitted for OPD more frequently than men, while the combination of OPD and pneumonia was a more frequent diagnosis amongst men (64%) (both $p < 0.001$). The condition was classified as being mild, moderate or serious/life threatening in 84, 163 and 90 patients, respectively. A higher proportion of patients with pneumonia were classified as being more severely ill (moderate, serious/life threatening) compared to those with OPD (82% versus 69%; $p < 0.05$). There was no significant association between the patient's age and the initial diagnosis ($p = 0.321$).

Recording of variables

Heart rate, body temperature and CRP were recorded in almost all patients – i.e. in 99%, 97% and 91%, respectively (see Table 2). Arterial blood gas concentrations were recorded in 78%, respiratory rate in 66%, PEF in 49%, and cutaneous oxygen saturation in 45% of patients. Arterial blood gas concentrations were recorded more frequently in men (84%) than in women (75%) ($p < 0.05$).

Measurement of PEF was performed in two-thirds of the patients with OPD, in approximately half of the patients with a combination diagnosis of OPD and pneumonia, but only in one-sixth of the patients with pneumonia ($p < 0.001$). Measurements of respiratory rate, PEF, arterial blood gas concentration and cutaneous oxygen saturation were all associated with the initial diagnosis ($p < 0.001$) – Table 2. Cutaneous oxygen saturation was more frequently recorded in patients with OPD and concurrent pneumonia than in

Measurement of physiological variables in the emergency room

Table 1. Age, sex and disease severity in 338 patients presenting with obstructive pulmonary disease, pneumonia or a combination of obstructive pulmonary disease and pneumonia.

	Obstructive pulmonary disease	Pneumonia	Obstructive pulmonary disease and pneumonia	All	p-value*
N	176	103	59	338	
Age, average (SD) ; years	63 (19)	63 (20)	66 (17)	64 (19)	0.321
Sex;					
Women; %	63	50	36	54	0.001
Men; %	37	50	64	46	
Disease severity;					
Mild; %	31	18	17	25	0.026
Moderate; %	42	52	62	48	
Serious/life treating; %	27	30	21	27	

*Pearson's Chi-square test

Table 2. Total number and percentage of patients in whom respiratory rate, heart rate, body temperature, peak expiratory flow, cutaneous oxygen saturation, arterial oxygen tension, arterial carbon dioxide tension and venous C-reactive protein were recorded by diagnosis.

	Obstructive pulmonary disease		Pneumonia		Obstructive pulmonary disease and pneumonia		In total		p-value*
	n	%	n	%	n	%	n	%	
N	176		103		59		338		
Heart rate (heart beats/min)	175	99	102	99	59	100	336	99	0.739
Body temperature (°C)	169	96	103	100	57	97	329	97	0.166
Respiratory rate (/min)	131	74	49	48	42	71	222	66	<0.001
Peak expiratory flow (PEF) (l/min)									
Men	45	69	9	18	21	55	75	48	<0.001
Women	73	66	7	14	9	43	89	49	
Cutaneous oxygen saturation (%)	90	51	29	28	33	56	152	45	<0.001
Arterial blood gases									
P _a O ₂ (kPa)	148	84	63	61	51	86	262	78	<0.001
P _a CO ₂ (kPa)	148	84	63	61	52	88	263	78	
C-reactive protein (mg/L)	156	89	97	94	54	92	307	91	0.019

*Pearson's Chi-square test

patients who only had pneumonia ($p < 0.001$).

Arterial blood gas concentrations were recorded less frequently in patients below the age of 45 (66%) than in patients aged 65 to 79 years (82%), and most frequently in patients older than 80 years (93%) ($p < 0.001$). Arterial blood gas concentrations were recorded in almost nine out of 10 patients with OPD or OPD and pneumonia, but only in two out of three patients with pneumonia ($p < 0.001$). CRP was recorded more frequently when the initial diagnosis was pneumonia than when the initial diagnosis was OPD ($p < 0.05$).

Peak expiratory flow and respiratory rate were far more frequently recorded in patients admitted to Haukeland University Hospital (58% and 72%, respectively) than in patients admitted to the other two hospitals (30% and 58%,

respectively; $p < 0.01$). CRP was more frequently recorded in Haugesund County Hospital and Haraldsplass Deaconal Hospital (94% of patients) than in Haukeland University Hospital (90%) ($p < 0.05$).

Multivariate analyses

In a multiple logistic regression adjusted for sex, age, diagnosis and disease severity we found that respiratory rate, PEF, cutaneous oxygen saturation and arterial blood gas concentrations were less frequently recorded in patients with pneumonia than in patients with OPD (Table 3). Cutaneous oxygen saturation and arterial blood gas concentrations were, as expected, more frequently measured in patients with serious or life threatening disease, whereas respiratory rate and PEF seemed to be recorded independently of disease

